FIREBIRD
BIOMOLECULAR SCIENCES, LLC

2026
Product Catalog

Nucleic acid analogs for:
- DNA sequencing
- Diagnostics
- SNP detection
- Synthetic Biology

www.firebirdbio.com

Contents

Reversible terminators 4
DNA sequencing, SNP identification, & *de novo* DNA synthesis

AEGIS™ 8
Artificially Expanded Genetic Information Systems
Orthogonality, cleanliness, and uniformity in nucleic acid analysis

SAMRS™ 17
Self-Avoiding Molecular Recognition Systems
Reducing primer dimers in nucleic acid amplification

SNAP-2 22
Identifying DNA and RNA molecules when the exact sequence is not known

Ordering 23

References 24

FIREBIRD BIOMOLECULAR SCIENCES, LLC

Firebird reagents and the origins of life



This catalog presents reagents and technologies to aid in the development of new biomedical innovations. Several of the products presented here were developed with the goal of better understanding the features of DNA and RNA that make them the carriers of genetic information for life on earth (Karalkar and Benner, 2018).

The image shown, left, illustrates the crystal structure of a double helix built from an expanded genetic alphabet of eight hachimoji building blocks, G (green), A (red), C (dark blue), T (yellow), B (aqua) P (blue), S (violet), and Z (orange). This hachimoji DNA forms the basis of Firebird. Adopted from Hoshika et al., 2019.

Reagents in this catalog are sold for research use only. Please inquire for diagnostic or other uses.

Reversible Terminators

Firebird has introduced the 3'-ONH_2 reversible terminator as an alternative to the larger 3'-OCH_2N_3 group (Hutter et al., 2010). The 3'-ONH_2 group is accepted by a variety of polymerases. After incorporation, further primer extension is blocked. The 3'-ONH_2 group is cleaved with buffered aqueous sodium nitrite to regenerate a standard 3'-OH.



3'-ONH_2 reversible terminators can be used in DNA sequencing (Hutter et al., 2010), Oligonucleotide synthesis (Jensen and Davis, 2018, Sarac and Hollenstein, 2019) and SNP analysis (Chen et al., 2010).

Polymerase variants have been developed that incorporate these terminators with improved efficiency over standard Taq DNA polymerase (Chen et al., 2010):

POL475-400	400 units	$132.00
POL475-1000	1000 units	$300.00

Reversible Terminators: Ready to Use, Untagged

Firebird offers “ready-to-use” un-tagged reversible terminators with a free 3'-ONH_2 group that can be directly incorporated by various enzymes without further manipulation.



Thymine
TONH2-171
5 µmoles $690.00
50 µmoles $ 4,830.00



Cytosine
CONH2-172
5 µmoles $863.00
50 µmoles $6,038.00



Adenine
AONH2-173
5 µmoles $863.00
50 µmoles $6,038.00



Guanine
GONH2-174
5 µmoles $863.00
50 µmoles $6,038.00



Inosine
IONH2-175
5 µmoles $863.00
50 µmoles $6,038.00

Please inquire for bulk pricing.

Reversible Terminators: diol-linked tags

Firebird offers reversible terminators with a 3'-ONH_2 (protected as the acetoxime) and a diol linker carrying a free amino group, to which a tag (fluorescent dye or other moiety) can be attached. The diol can be rapidly cleaved with dilute aqueous periodate.
For enzymatic incorporation, the oxime must be deprotected to the free 3'-ONH_2 before use, which can be achieved in situ with buffered aqueous methoxylamine.



Thymine
TONH2-DT
1 µmole $2,013.00



Cytosine
CONH2-DT
1 µmole $2,185.00



Adenine
7c8n-AONH2-DT
1 µmole $2,530.00



Guanine
7c8n-GONH2-DT
1 µmole $2,645.00

Please inquire for bulk pricing.

Not available for sequencing applications on certain machines and in certain jurisdictions. Please inquire.

Reversible Terminators: disulfide-linked tags

Firebird offers reversible terminators with a 3'-ONH_2 (protected as the acetoxime) and a disulfide linker carrying a free amino group, to which a tag (fluorescent dye or other moiety) can be attached. The disulfide can be rapidly cleaved with phosphine or thiol reagents. For enzymatic incorporation, the oxime must be deprotected to the free 3'-ONH_2 before use, which can be achieved in situ with buffered aqueous methoxylamine.



Thymine
TONH2-ST
1 µmole $ 2,013.00



Cytosine
CONH2-ST
1 µmole $ 2,300.00



Adenine
7c8n-AONH2-ST
1 µmole $ 2,760.00



Guanine
7c8n-GONH2-ST
1 µmole $ 2,875.00

Please inquire for bulk pricing.

Not available for sequencing applications on certain machines and in certain jurisdictions. Please inquire.

AEGIS

Artificially Expanded Genetic Information Systems

Firebird has created different heterocycles in order to implement additional hydrogen bonding patterns for the Z:P and S:B base pairs as shown below. These base pairs are "orthogonal" to C:G and A:T (Benner, 2004; Hoshika et al., 2019; Sefah et al., 2014; Yang et al., 2006).











FIREBIRD BIOMOLECULAR SCIENCES, LLC

Using AEGIS:
For Ultra-clean Nested PCR

The Z:P pair is retained during PCR amplification, even when present as consecutive base pairs (Yang et al., 2011).





AEGIS-nested primers suppress noise in multiplex PCR (Yang et al., 2010) and support 22-plex PCR to detect RNA viruses (Glushakova et al., 2015a).





Using AEGIS:
Assembling Long DNA Constructs

Adding AEGIS nucleotides to single-stranded oligos increases their information density. This allows clean and rapid hybridization of single stranded DNA unobstructed by hairpins, wandering strands, and non-canonical interactions. Then, using Firebird's "transliteration" technology, the AEGIS nucleotides are cleanly replaced by standard nucleotides giving an entirely natural gene (Bradley and Benner, 2014; Merritt et al., 2014).

An example of this assembly is shown below: S and B residues in the tails of the primers guide specific alignment of the unique sequences at the ends of the primers. The S:B basepair is converted to a T:A basepair during PCR amplification.



Using AEGIS:
Molecular Beacons

Adding AEGIS nucleotides to the stems of molecular beacons prevents their being opened by adventitious DNA and RNA in a complex biological sample. This allows beacons to deliver signals in complex biological samples with significantly lower noise (Sheng et al., 2008).



Firebird offers custom-synthesized beacons with a wide range of fluorescent dyes and quenchers. Please inquire.

AEGIS:

Increasing Aptamer Diversity

The dP and dZ bases increase the chemical repertoire of nuleic acids for aptamer selections (Biondi et al., 2016; Sefah et al., 2014; Zhang et al., 2015, 2016). Of special note is the NO_2 group present on dZ.

The figure below shows one application of AEGIS for the selection of aptamers specific to a cell surface protein on a cancer cell (Zhang et al., 2016).



AEGIS:
dZ and dP Phosphoramidites

dZ Phosphoramidite



Hydrogen bonding pattern: Z
Sugar: 2'-Deoxyribose
Heterocycle: Nitropyridine
Linear Formula: $C_{48}H_{55}N_6O_{11}P$
Mol Weight: 922.96

Cat. No. dZ-PA-101
100 mg $ 828.00
1 gram $6,62400

dP Phosphoramidite



Hydrogen bonding pattern: P
Sugar: 2'-Deoxyribose
Heterocycle: Imidazotriazine
Linear Formula: $C_{49}H_{65}N_8O_7P$
Mol Weight: 909.08

Cat. No. dP-PA-102
100 mg $ 978.00
1 gram $7,820.00

(Yang et al., 2006)
Please inquire for bulk pricing

Oligonucleotides are available that contain dZ and dP.
Please inquire for availability of ribonucleoside derivatives.

AEGIS:

dZ and dP Triphosphates

dZ Triphosphate



Hydrogen bonding pattern: Z
Sugar: 2'-deoxyribose
Heterocycle: Nitropyridine
Linear Formula: $C_{10}H_{16}N_3O_{15}P_3$
Mol Weight: 511.17

Cat. No. dZTP-101
1 µmole $ 460.00
5 µmoles $ 1,840.00

dP Triphosphate



Hydrogen bonding pattern: P
Sugar: 2'-deoxyribose
Heterocycle: Imidazotriazine
Linear Formula: $C_{10}H_{16}N_5O_{13}P_3$
Mol Weight: 493.15

Cat. No. dPTP-201
1 µmole $ 483.00
5 µmoles $ 1,932.00

(Yang et al., 2007)
Please inquire for bulk pricing.

Please inquire for availability of ribonucleoside derivatives.

AEGIS:
dS and dB Phosphoramidites

dS Phosphoramidite



Hydrogen bonding pattern: S
Sugar: 2'-Deoxyribose
Heterocycle: Pseudo-C
Linear Formula: $C_{47}H_{54}N_5O_8P$
Mol Weight : 847.93

Cat. No. dS-PA-104S

100 mg	$ 828.00
1 gram	$6,62400

dB Phosphoramidite



Hydrogen bonding pattern: B
Sugar: 2'-Deoxyribose
Heterocycle: Purine
Linear Formula: $C_{62}H_{79}N_8O_8P$
Mol Weight: 1095.31

Cat. No. dB-PA-103P

100 mg	$ 720.00
1 gram	$5,760.00

Please inquire for bulk pricing.

Oligonucleotides are available that incorporate dS and dB.
Please inquire for availability of ribonucleoside derivatives.

AEGIS:

dS and dB Triphosphates

dS Triphosphate



Hydrogen bonding pattern: S
Sugar: 2'-deoxyribose
Heterocycle: Pseudo-C
Linear Formula: $C_{10}H_{18}N_3O_{13}P_3$
Mol Weight: 481.18

Cat. No. dSTP-401S
1 µmole $ 460.00
5 µmoles $ 1,840.00

dB Triphosphate



Hydrogen bonding pattern: B
Sugar: 2'-deoxyribose
Heterocycle: Purine
Linear Formula: $C_{10}H_{16}N_5O_{13}P_3$
Mol Weight: 507.18

Cat. No. dBTP-301P
1 µmole $ 460.00
5 µmoles $ 1,840.00

Please inquire for bulk pricing.

Please inquire for availability of ribonucleoside derivatives.

FIREBIRD BIOMOLECULAR SCIENCES, LLC

SAMRS™

Self Avoiding Molecular Recognition System

Nucleic acid amplification reactions commonly suffer from the formation of off-target amplification products. This problem increases with the number of primers in the reaction. It mainly occurs because of primer-primer interactions. Thus, selectively removing hydrogen bonding units from the primer bases results in *self-avoiding* DNA primers that can be added to enable and improve multiplexed PCR (Hoshika et al., 2010).

As shown below, SAMRS bases, indicated with a *, can base pair with standard bases (in the target or amplicon) but not with other SAMRS bases.



FIREBIRD BIOMOLECULAR SCIENCES, LLC

SAMRS:

Phosphoramidites

g* phosphoramidite

1 g	$ 240
10 g	$ 2,040

a* phosphoramidite

1 g	$ 753
10 g	$ 6,141

c* phosphoramidite

1 g	$ 453
10 g	$ 3,695

t* phosphoramidite

1 g	$ 1,208
10 g	$ 9,820

Oligonucleotides are available that incorporate the SAMRS bases.

SAMRS can be combined with AEGIS in oligos to get highly multiplexed and highly clean PCR, as well as isothermal amplification. Please inquire.

FIREBIRD BIOMOLECULAR SCIENCES, LLC

SAMRS:

Primer & Probe Sets - *SARS-CoV-2*

SAMRS-modified primer and probe sets for the identification of SARS-CoV-2 (2019-nCoV).

All sets include individual primers and probes as specified, HPLC purified and delivered in IDTE (1X TE buffer) pH 7.5 at 100µM. Available in normalized deliverables to the amounts indicated below and shipped ambient.

Primer/Probe	Quantity (nmol)	Price (USD)
nCOV_N1 Forward SAMRS Primer	50 nmol	$200.00
nCOV_N1 Reverse SAMRS Primer	50 nmol	$200.00
nCOV_N1 (FAM) Probe	25 nmol	$350.00
nCOV_N2 Forward SAMRS Primer	50 nmol	$200.00
nCOV_N2 Reverse SAMRS Primer	50 nmol	$200.00
nCOV_N2 (FAM) Probe	25 nmol	$350.00
nCOV_N3 Forward SAMRS Primer	50 nmol	$200.00
nCOV_N3 Reverse SAMRS Primer	50 nmol	$200.00
nCOV_N3 (FAM) Probe	25 nmol	$350.00
RNase P Forward SAMRS Primer	50 nmol	$200.00
RNase P Reverse SAMRS Primer	50 nmol	$200.00
RNase P (FAM) Probe	25 nmol	$350.00
RNase P (Cy 5) Probe	25 nmol	$700.00
nCOV_N1 Forward SAMRS Primer	100 nmol	$300.00
nCOV_N1 Reverse SAMRS Primer	100 nmol	$300.00
nCOV_N1 (FAM) Probe	50 nmol	$530.00
nCOV_N2 Forward SAMRS Primer	100 nmol	$300.00
nCOV_N2 Reverse SAMRS Primer	100 nmol	$300.00
nCOV_N2 (FAM) Probe	50 nmol	$530.00
nCOV_N3 Forward SAMRS Primer	100 nmol	$300.00
nCOV_N3 Reverse SAMRS Primer	100 nmol	$300.00
nCOV_N3 (FAM) Probe	50 nmol	$530.00
RNase P Forward SAMRS Primer	100 nmol	$300.00
RNase P Reverse SAMRS Primer	100 nmol	$300.00
RNase P (FAM) Probe	50 nmol	$530.00
RNase P (Cy 5) Probe	50 nmol	$1,000.00
Complete Primer & Probe Set		**$9,220.00**

Primers & Probes are available individually or as a complete set.
SAMRS-modified Primers and probes are manufactured in a template-free environment and certified template-free to cycle 45 by NTC testing.
For Research Use Only.

SAMRS:

Primer & Probe Sets - *SARS-CoV-2 and Influenza*

SAMRS-modified primer and probe sets for the identification of SARS-CoV-2 (2019-nCoV) and Influenza (InfA and InfB).

All sets include individual primers and probes as specified, HPLC purified and delivered in IDTE (1X TE buffer) pH 7.5 at 100µM. Available in normalized deliverables to the amounts indicated below and shipped ambient.

Primer/Probe	Quantity (nmol)	Price
InfA Forward 1 SAMRS Primer	100 nmol	$300.00
InfA Forward 2 SAMRS Primer	100 nmol	$300.00
InfA Reverse 1 SAMRS Primer	100 nmol	$300.00
InfA Reverse 2 SAMRS Primer	100 nmol	$300.00
InfA (FAM) Probe	25 nmol	$350.00
InfB Forward SAMRS Primer	100 nmol	$300.00
InfB Reverse SAMRS Primer	100 nmol	$300.00
InfB (YAK) Probe	25 nmol	$450.00
SC2 Forward SAMRS Primer	100 nmol	$300.00
SC2 Reverse SAMRS Primer	100 nmol	$300.00
SC2 (TexRd) Probe	25 nmol	$700.00
RNase P Forward SAMRS Primer	100 nmol	$300.00
RNase P Reverse SAMRS Primer	100 nmol	$300.00
RNase P (Cy 5) Probe	25 nmol	$700.00
Complete Primer & Probe Set		**$5,200.00**

Primers & Probes are available individually or as a complete set.

SAMRS-modified Primers and probes are manufactured in a template-free environment and certified template-free to cycle 45 by NTC testing.

For Research Use Only.

SAMRS:
Design Rules

General rules for placement of SAMRS bases into primers used in PCR and other amplification techniques. Upper case letters indicate the standard base, lower case letters indicate the SAMRS base.

1. Recommended lengths are 20-35 nts. SAMRS bases should be utilized in the first 4, (up to 8), positions at the 3' end of the oligonucleotide but not in the very first 3' base.

2. Between two and four SAMRS bases should be used per oligonucleotide, with two or three SAMRS modifications being preferred.

3. The SAMRS t base has not demonstrated as much reduction in primer dimer formation so it is preferable to substitute a, g or c rather than t when given a choice.

4. Avoid using SAMRS bases in a string of three and four consecutive identical bases. For example: ggg, ccc, aaa, ttt, gggg, cccc, aaaa, and tttt are to be avoided.

5. Separate SAMRS bases with standard bases, for example: gGg, cCc, aAa, and tTt.

6. If the 3'-end of an oligo is standard A or T, the second base should also be a standard base, followed by two or three SAMRS modifications.

(Benner et al., 2015; Glushakova et al., 2015a, 2015b; Hoshika et al., 2010; Sharma et al., 2014; Yang et al., 2015 and unpublished data)

As with all primer designs, Firebird makes no guarantee as to the performance of oligonucleotides made by following these rules.

SNAP2™ Oligonucleotides: Priming with High Specificity

Specific DNA priming requires oligonucleotides with complementarity of ≥16 nucleotides but the ability to discriminate against single mismatches. Dynamic assembly of a primer on a template can provide this (Leal et al., 2006).



Controlled pore glass with aldehyde precursor
SNAP2T_HJK001
100 mg $600.00

Firebird sells phosphoramidites, oligos and libraries with SNAP2 ends. Please inquire.

SNAP2 primers are not to be confused with Snap-Tag® and other registered trademarks of New England BioLabs.

Purchasing, Shipping and Handling

Contact

Orders
By email: orders@firebirdbio.com
By telephone: (386) 418-0347
9:00 - 5:00 Eastern Time

Technical inquiries
support@firebirdbio.com

Reagents and enzymes are shipped at the customer's expense, on dry ice as appropriate.

References

Benner, S.A. (2004). Chemistry. Redesigning genetics. Science 306, 625–626.

Benner, S.A., Kim, H.-J., Merritt, K.B., Yang, Z., McLendon, D.C., Hoshika, S., and Hutter, D. (2015). Next-generation DNA in pathogen detection, surveillance, and CLIA-waivable diagnostics. SPIE Digital Library 9490, 94900K-94900K – 6.

Biondi, E., Lane, J.D., Das, D., Dasgupta, S., Piccirilli, J.A., Hoshika, S., Bradley, K.M., Krantz, B.A., and Benner, S.A. (2016). Laboratory evolution of artificially expanded DNA gives redesignable aptamers that target the toxic form of anthrax protective antigen. Nucleic Acids Res. 44, 9565–9577, PMCID: PMC5175368.

Bradley, K.M., and Benner, S.A. (2014). OligArch: A software tool to allow artificially expanded genetic information systems (AEGIS) to guide the autonomous self-assembly of long DNA constructs from multiple DNA single strands. Beilstein J Org Chem 10, 1826–1833, PMCID: PMC4142867.

Chen, F., Gaucher, E.A., Leal, N.A., Hutter, D., Havemann, S.A., Govindarajan, S., Ortlund, E.A., and Benner, S.A. (2010). Reconstructed evolutionary adaptive paths give polymerases accepting reversible terminators for sequencing and SNP detection. Proc. Natl. Acad. Sci. U.S.A. 107, 1948–1953, PMCID: PMC2804741.

Glushakova, L.G., Bradley, A., Bradley, K.M., Alto, B.W., Hoshika, S., Hutter, D., Sharma, N., Yang, Z., Kim, M.-J., and Benner, S.A. (2015a). High-throughput multiplexed xMAP Luminex array panel for detection of twenty two medically important mosquito-borne arboviruses based on innovations in synthetic biology. J. Virol. Methods 214, 60–74, PMCID: PMC4485418.

Glushakova, L.G., Sharma, N., Hoshika, S., Bradley, A.C., Bradley, K.M., Yang, Z., and Benner, S.A. (2015b). Detecting respiratory viral RNA using expanded genetic alphabets and self-avoiding DNA. Anal. Biochem. 489, 62–72, PMCID: PMC4733849.

Hoshika, S., Chen, F., Leal, N.A., and Benner, S.A. (2010). Artificial genetic systems: self-avoiding DNA in PCR and multiplexed PCR. Angew. Chem. Int. Ed. Engl. 49, 5554–5557, PMCID: PMC6027612.

Hoshika, S., Leal, N.A., Kim, M.-J., Kim, M.-S., Karalkar, N.B., Kim, H.-J., Bates, A.M., Watkins, N.E., SantaLucia, H.A., Meyer, A.J., et al. (2019). Hachimoji DNA and RNA: A genetic system with eight building blocks. Science 363, 884–887, PMCID: PMC6413494.

FIREBIRD BIOMOLECULAR SCIENCES, LLC

References

Hutter, D., Kim, M.-J., Karalkar, N., Leal, N.A., Chen, F., Guggenheim, E., Visalakshi, V., Olejnik, J., Gordon, S., and Benner, S.A. (2010). Labeled nucleoside triphosphates with reversibly terminating aminoalkoxyl groups. Nucleosides Nucleotides Nucleic Acids 29, 879–895, PMCID: PMC3858015.

Jensen, M.A., and Davis, R.W. (2018). Template-Independent Enzymatic Oligonucleotide Synthesis (TiEOS): Its History, Prospects, and Challenges. Biochemistry 57, 1821–1832.

Karalkar, N.B., and Benner, S.A. (2018). The challenge of synthetic biology. Synthetic Darwinism and the aperiodic crystal structure. Curr Opin Chem Biol 46, 188–195.

Leal, N.A., Sukeda, M., and Benner, S.A. (2006). Dynamic assembly of primers on nucleic acid templates. Nucleic Acids Res. 34, 4702–4710, PMCID: PMC1635275.

Merritt, K.K., Bradley, K.M., Hutter, D., Matsuura, M.F., Rowold, D.J., and Benner, S.A. (2014). Autonomous assembly of synthetic oligonucleotides built from an expanded DNA alphabet. Total synthesis of a gene encoding kanamycin resistance. Beilstein J Org Chem 10, 2348–2360, PMCID: PMC4222377.

Sarac, I., and Hollenstein, M. (2019). Terminal Deoxynucleotidyl Transferase in the Synthesis and Modification of Nucleic Acids. Chembiochem 20, 860–871.

Sefah, K., Yang, Z., Bradley, K.M., Hoshika, S., Jiménez, E., Zhang, L., Zhu, G., Shanker, S., Yu, F., Turek, D., et al. (2014). In vitro selection with artificial expanded genetic information systems. Proc. Natl. Acad. Sci. U.S.A. 111, 1449–1454, PMCID: PMC3910645.

Sharma, N., Hoshika, S., Hutter, D., Bradley, K.M., and Benner, S.A. (2014). Recombinase-based isothermal amplification of nucleic acids with self-avoiding molecular recognition systems (SAMRS). Chembiochem 15, 2268–2274.

Sheng, P., Yang, Z., Kim, Y., Wu, Y., Tan, W., and Benner, S.A. (2008). Design of a novel molecular beacon: modification of the stem with artificially genetic alphabet. Chem. Commun. (Camb.) 5128–5130, PMCID: PMC2763601.

FIREBIRD BIOMOLECULAR SCIENCES, LLC

References

Yang, Z., Hutter, D., Sheng, P., Sismour, A.M., and Benner, S.A. (2006). Artificially expanded genetic information system: a new base pair with an alternative hydrogen bonding pattern. Nucleic Acids Res. 34, 6095–6101, PMCID: PMC1635279.

Yang, Z., Sismour, A.M., Sheng, P., Puskar, N.L., and Benner, S.A. (2007). Enzymatic incorporation of a third nucleobase pair. Nucleic Acids Res. 35, 4238–4249, PMCID: PMC1934989.

Yang, Z., Chen, F., Chamberlin, S.G., and Benner, S.A. (2010). Expanded genetic alphabets in the polymerase chain reaction. Angew. Chem. Int. Ed. Engl. 49, 177–180, PMCID: PMC3155763.

Yang, Z., Chen, F., Alvarado, J.B., and Benner, S.A. (2011). Amplification, mutation, and sequencing of a six-letter synthetic genetic system. J. Am. Chem. Soc. 133, 15105–15112, PMCID: PMC3427765.

Yang, Z., McLendon, C., Hutter, D., Bradley, K.M., Hoshika, S., Frye, C.B., and Benner, S.A. (2015). Helicase-Dependent Isothermal Amplification of DNA and RNA by Using Self-Avoiding Molecular Recognition Systems. Chembiochem 16, 1365–1370, PMCID: PMC4489552.

Yang, Z., Kim, H.-J., Le, J.T., McLendon, C., Bradley, K.M., Kim, M.-S., Hutter, D., Hoshika, S., Yaren, O., and Benner, S.A. (2018). Nucleoside analogs to manage sequence divergence in nucleic acid amplification and SNP detection. Nucleic Acids Res. 46, 5902–5910, PMCID: PMC6159519.

Zhang, L., Yang, Z., Sefah, K., Bradley, K.M., Hoshika, S., Kim, M.-J., Kim, H.-J., Zhu, G., Jiménez, E., Cansiz, S., et al. (2015). Evolution of functional six-nucleotide DNA. J. Am. Chem. Soc. 137, 6734–6737, PMCID: PMC4500535.

Zhang, L., Yang, Z., Le Trinh, T., Teng, I.-T., Wang, S., Bradley, K.M., Hoshika, S., Wu, Q., Cansiz, S., Rowold, D.J., et al. (2016). Aptamers against Cells Overexpressing Glypican 3 from Expanded Genetic Systems Combined with Cell Engineering and Laboratory Evolution. Angew. Chem. Int. Ed. Engl. 55, 12372–12375, PMCID: PMC5554412.

FIREBIRD BIOMOLECULAR SCIENCES, LLC